

# On Generalized Seventh Order Pell And Pell-Like Sequences

Nasir Ahmad, Leena Prasher

Department of Mathematics CT University, Ferozpur Road, Ludhiana, Punjab, India.

**Article Info**

Article history:

Received 29 June 2021

Received in revised form

28 November 2021

Accepted 5 December 2021

Available online 15 December 2021

**Keywords** Generating function, Pell Sequence, seventh order, Binet Formulae.

**Abstract:** The generalizations of Fibonacci sequence have wide range of properties and applications in every field of science and hybrid science. The Fibonacci sequence has been generalized in numerous ways either with the same initial conditions or by altering the recurrence relation and vis a vis In this regard, we attempted to consummate all relevant and available literature in order to provide readers with a solid foundation for further scientific research for higher order Pell-like sequences. In this paper, we present some results on the generalized Pell sequences and Pell-like sequences of order seven. The generating function, Binet Formula and linear sum for Pell, Pell-Lucas and modified Pell sequences of order seven will be investigated and results on them. Also some well-known identities for order seven will be presented for the same.

**1. Introduction**

In this paper, we introduce the generalized Pell sequences of order seven and we investigate Pell-like sequences; Pell-Lucas and modified Pell sequences of order seven. The Pell sequence  $\{P_n\}$  is defined recursively by

$$P_{n+2} = 2P_{n+1} + P_n; n \geq 0; P_0 = 0, P_1 = 1; \tag{1.1}$$

First few terms of Pell sequence are; 0, 1, 2, 5, 12, 29, 70, 169... OEIS: A000129, [13].

Several authors and researchers have presented and evaluated Pell sequence in different aspects [1-19]. The second order recurrence relations of Pell numbers have been generalized. Generalized Pell numbers and their properties have been studied by various authors [20-23].

This paper can provide new holistic approach for further expansion of research for the higher order generalized Pell sequences and Pell-like sequences not only in Number theory but in engineering, computer sciences and combinatorics. The generalization of Pell sequence for order third, fourth and fifth are presented by Soykan [14-20].

A generalized seventh order Pell sequence  $\{H_m\}_{m \geq 0} = \{H_m(H_{j=0,1,\dots,6}; \lambda_{r=1,2,\dots,7})_{m \geq 0} \in \mathbb{R}$  with the conditions  $\lambda_1 = 2, \lambda_2 = 1, \lambda_3 = \lambda_4 = \lambda_5 = \lambda_6 = \lambda_7$  is defined by the seventh-order recurrence relations as

$$H_m = 2H_{m-1} + \sum_{j=2}^7 H_{m-j} \tag{1.2}$$

The characteristic equation of is

$$y^7 - 2y^6 - \sum_{j=0}^5 y^j = 0 \tag{1.3}$$

Where are the roots of the above characteristic equation (1.3) such

$$\sum_{j=1}^7 \alpha_j = 2 \quad \prod_{j=1}^7 \alpha_j = 1 \tag{1.4}$$

The first few generalized seventh order Pell numbers for are given as;

$$H_0, H_1, H_2, H_3, H_4, H_5, H_6, 2H_6 + \sum_{j=0}^5 H_j, 2H_7 + \sum_{j=1}^6 H_j, 2H_8 + \sum_{j=2}^7 H_j, 2H_9 + \sum_{j=3}^8 H_j \tag{1.5}$$

**Corresponding Author,**

**E-mail address:**

Leena.17064@ctuniversity.in

All rights reserved: <http://www.ijari.org>

syeednasir25@gmail.com;

Now let  $\{P_m^*\}_{m \geq 0}$ ,  $\{Q_m^*\}_{m \geq 0}$  and  $\{E_m^*\}_{m \geq 0}$  be the Pell-like sequence, Pell-Lucas sequence and modified Pell sequence from  $H_m$  of order seven defined by the recurrence relations for each as;

$$P_{m+7}^* = 2P_{m+5}^* + \sum_{j=0, j \neq 5}^6 P_{m+j}^*, Q_{m+7}^* = 2Q_{m+5}^* + \sum_{j=0, j \neq 5}^6 Q_{m+j}^* \\ E_{m+7}^* = 2E_{m+5}^* + \sum_{j=0, j \neq 5}^6 E_{m+j}^* \tag{1.6}$$

With the first few terms of each sequence as and

With the first few terms of each sequence as and

$$P_0^* = 0, P_1^* = 1, P_2^* = 2, P_3^* = 5, \dots, Q_0^* = 4, Q_1^* = 2, Q_2^* = 6, Q_3^* = 17, \dots \\ E_0^* = 0, E_1^* = 1, E_2^* = 1, E_3^* = 3, \dots \tag{1.7}$$

The OEIS database does not contain sequences of  $\{P_m^*\}_{m \geq 0}$ ,  $\{Q_m^*\}_{m \geq 0}$  and  $\{E_m^*\}_{m \geq 0}$  [13], yet.

**2. Ordinary generating functions of  $\{H_m\}_{m \geq 0}$  Lemma**

Let  $fH_m(y) = \sum_{m=0}^{\infty} H_m y^m$  is the ordinary generating function

of the generalized seventh-order Pell sequence  $\{H_m\}$  : Then,

$\sum_{m=0}^{\infty} H_m y^m$  is given by

$$\sum_{m=0}^{\infty} H_m y^m = \frac{H_0 + (H_1 - 2H_0)y + (H_2 - 2H_1 - H_0)y^2 + \dots + (H_6 - 2H_5 - H_4 - H_3 - H_2 - H_1 - H_0)y^6}{1 - 2y - \sum_{m=2}^7 y^m}$$

Proof. By definition of generalized seventh-order Pell numbers and subtracting  $y^j f(y); j = 1, 2, \dots, 7$  from  $f(y)$  respectively we have

$$(1 - 2y - y^2 - y^3 - y^4 - y^5 - y^6 - y^7) fH_m(y) = \sum_{m=0}^{\infty} H_m y^m - 2y \sum_{m=0}^{\infty} H_m y^m - y^2 \sum_{m=0}^{\infty} H_m y^m - \dots - y^7 \sum_{m=0}^{\infty} H_m y^m$$

$$\begin{aligned}
 &= \sum_{m=0}^{\infty} H_m y^m - 2 \sum_{m=0}^{\infty} H_m y^{m+1} - y \sum_{m=0}^{\infty} H_m y^{m+2} - \dots - y \sum_{m=0}^{\infty} H_m y^{m+7} \\
 &= \sum_{m=0}^{\infty} H_m y^m - 2 \sum_{m=1}^{\infty} H_{m-1} y^{m+1} - y \sum_{m=2}^{\infty} H_{m-2} y^{m+2} - \dots - y \sum_{m=7}^{\infty} H_{m-7} y^{m+7} \\
 &= \left( \sum_{j=0}^6 H_j y^j \right) - 2 \left( \sum_{j=0}^6 H_j y^j \right)
 \end{aligned}$$

By rearrangement of terms, result holds.

**2.1 Corollary 1.**

The generated functions for Pell, Pell-Lucas and modified Pell sequences of order seven are respectively as;

$$\begin{aligned}
 \sum_{m=0}^{\infty} P_m^* y^m &= \frac{y}{1 - 2y - \sum_{i=2}^7 y^i} \\
 \sum_{m=0}^{\infty} Q_m^* y^m &= \frac{7 - 12y - 5y^2 - 4y^3 - 3y^4 - 2y^5 - y^6}{1 - 2y - \sum_{i=2}^7 y^i} \quad \text{and} \\
 \sum_{m=0}^{\infty} E_m^* y^m &= \frac{y - y^2}{1 - 2y - \sum_{i=2}^7 y^i}
 \end{aligned}$$

**3. Obtaining binet formula from generating function**

**3.1 Theorem 1**

Generalized Binet formula for  $\{H_m\}$  of Pell numbers for order seven

$$\{H_m\} = \frac{d_1 \alpha_1^m}{\prod_{k=2}^7 (\alpha_1 - \alpha_k)} + \frac{d_2 \alpha_2^m}{\prod_{k=1, k \neq 2}^7 (\alpha_2 - \alpha_k)} + \dots + \frac{d_7 \alpha_7^m}{\prod_{k=1}^6 (\alpha_7 - \alpha_k)} \tag{3.1}$$

Proof. Let

$$h(y) = 1 - 2y - y^2 - y^3 - y^4 - y^5 - y^7 - y^6 - y^7, \text{ then}$$

for some  $\alpha_i; i = 1, 2, \dots, 7$  we have

$$h\left(\frac{1}{y}\right) = 1 - \frac{2}{y} - \frac{1}{y^2} - \frac{1}{y^3} - \frac{1}{y^4} - \frac{1}{y^5} - \frac{1}{y^6} - \frac{1}{y^7} = 0$$

Hence  $\frac{1}{\alpha_i}, i = 1, 2, \dots, 7$  are the roots of  $h(y)$ : This gives

$\alpha_i; i = 1, 2, \dots, 7$  as the roots of

$$\begin{aligned}
 h(y) &= 1 - 2y - y^2 - y^3 - y^4 - y^5 - y^6 - y^7 \\
 \Rightarrow y^7 - y^6 - y^5 - y^4 - y^3 - 2y^2 - y - 1 &= 0
 \end{aligned}$$

Now, by (2.1) and (3.2), it follows that

$$\sum_{m=0}^{\infty} H_m y^m = \frac{H_0 + (H_1 - 2H_0)y + (H_2 - 2H_1 - H_0)y^2 + \dots + (H_6 - 2H_5 - H_4 - H_3 - H_2 - H_1 - H_0)y^6}{(1 - \alpha_1 y)(1 - \alpha_2 y)(1 - \alpha_3 y)(1 - \alpha_4 y)(1 - \alpha_5 y)(1 - \alpha_6 y)(1 - \alpha_7 y)}$$

Now

$$H_0 + (H_1 - 2H_0)y + (H_2 - 2H_1 - H_0)y^2 + \dots + (H_6 - 2H_5 - H_4 - H_3 - H_2 - H_1 - H_0)y^6$$

$$\begin{aligned}
 &= U_1 \prod_{i=2}^7 (1 - \alpha_i y) + U_2 \prod_{i=2, j \neq 2}^7 (1 - \alpha_i y) + U_3 \prod_{i=1, i \neq 3}^7 (1 - \alpha_i y) + U_4 \prod_{i=1, i \neq 4}^7 (1 - \alpha_i y) + U_5 \prod_{i=1, i \neq 5}^7 (1 - \alpha_i y) + \\
 &U_6 \prod_{i=1, i \neq 6}^7 (1 - \alpha_i y) + U_7 \prod_{i=1}^6 (1 - \alpha_i y)
 \end{aligned}$$

If consider  $y = \frac{1}{\alpha_1}$  we have

$$H_0 + (H_1 - 2H_0)\frac{1}{\alpha_1} + (H_2 - 2H_1 - H_0)\frac{1}{\alpha_1^2} + \dots + (H_6 - 2H_5 - H_4 - H_3 - H_2 - H_1 - H_0)\frac{1}{\alpha_1^6}$$

$$\begin{aligned}
 &= U_1 \left(1 - \frac{\alpha_2}{\alpha_1}\right) \left(1 - \frac{\alpha_3}{\alpha_1}\right) \left(1 - \frac{\alpha_4}{\alpha_1}\right) \left(1 - \frac{\alpha_5}{\alpha_1}\right) \left(1 - \frac{\alpha_6}{\alpha_1}\right) \left(1 - \frac{\alpha_7}{\alpha_1}\right) \\
 \Rightarrow U_1 &= \frac{\alpha_1^6 \left[ H_0 + (H_1 - 2H_0)\frac{1}{\alpha_1} + (H_2 - 2H_1 - H_0)\frac{1}{\alpha_1^2} + \dots + (H_6 - 2H_5 - H_4 - H_3 - H_2 - H_1 - H_0)\frac{1}{\alpha_1^6} \right]}{(\alpha_1 - \alpha_2)(\alpha_1 - \alpha_3)(\alpha_1 - \alpha_4)(\alpha_1 - \alpha_5)(\alpha_1 - \alpha_6)(\alpha_1 - \alpha_7)}
 \end{aligned}$$

$$\Rightarrow U_1 = \frac{d_1}{\prod_{i=2}^7 (\alpha_1 - \alpha_i)}$$

Similarly, we can obtain for  $U_k; k = 1, 2, \dots, 7$

$$\Rightarrow U_k = \frac{d_k}{\prod_{\substack{k, i=1 \\ k \neq i}}^7 (\alpha_k - \alpha_i)}$$

Thus (3.3) can be written as

$$\begin{aligned}
 \sum_{m=0}^{\infty} H_m y^m &= U_1 (1 - \alpha_1 y)^{-1} + U_2 (1 - \alpha_2 y)^{-1} + \dots + U_7 (1 - \alpha_7 y)^{-1} \\
 &= U_1 \sum_{m=0}^{\infty} \alpha_1^m y^m + U_2 \sum_{m=0}^{\infty} \alpha_2^m y^m + \dots + U_7 \sum_{m=0}^{\infty} \alpha_7^m y^m \\
 &= \sum_{m=0}^{\infty} (U_1 \alpha_1^m + U_2 \alpha_2^m + \dots + U_7 \alpha_7^m) y^m \\
 \Rightarrow H_m &= U_1 \alpha_1^m + U_2 \alpha_2^m + U_3 \alpha_3^m + \dots + U_7 \alpha_7^m
 \end{aligned}$$

**3.2 Corollary 2.**

Similarly, for seventh-order of Pell, Pell-Lucas and modified Pell sequences are as;

$$P_m^* = \sum_{k=1}^7 \frac{\alpha_k^{n+5}}{\prod_{\substack{k, i=1 \\ k \neq i}}^7 (\alpha_k - \alpha_i)}, \quad Q_m^* = \sum_{k=1}^7 \alpha_k^m \quad \text{and}$$

$$E_m^* = \sum_{k=1}^7 \frac{(\alpha_k - 1) \alpha_k^{n+4}}{\prod_{\substack{k, i=1 \\ k \neq i}}^7 (\alpha_k - \alpha_i)} \text{ respectively.}$$

**3.3 Corollary 3.**

The Binet formula of generalized seventh order Pell numbers can be represented as

$$H_m = \sum_{k=1}^7 \frac{\alpha_k d_k \alpha_k^m}{2\alpha_k^6 + 2\alpha_k^5 + 3\alpha_k^4 + 4\alpha_k^3 + 5\alpha_k^2 + 6\alpha_k + 7}$$

**3.4 Corollary 4**

$$\frac{1}{(\alpha_1 - \alpha_2)(\alpha_1 - \alpha_3)(\alpha_1 - \alpha_4)(\alpha_1 - \alpha_5)(\alpha_1 - \alpha_6)(\alpha_1 - \alpha_7)} = \frac{\alpha_1}{2\alpha_1^6 + 2\alpha_1^5 + 3\alpha_1^4 + 4\alpha_1^3 + 5\alpha_1^2 + 6\alpha_1 + 7}$$

**3.5 Corollary 5**

Similarly, we have for  $\alpha_i; i = 1, 2, \dots, 7$  we have

$$\frac{1}{\prod_{\substack{k, j=1 \\ k \neq j}}^7 (\alpha_k - \alpha_j)} = \frac{\alpha_k}{2\alpha_k^6 + 2\alpha_k^5 + 3\alpha_k^4 + 4\alpha_k^3 + 5\alpha_k^2 + 6\alpha_k + 7}$$

**Corollary 6.**

Binet formulas for Pell, Pell-Lucas, and modified Pell sequences of order seven as;

$$P_m^* = \sum_{k=1}^7 \frac{\alpha_k^{m+6}}{2\alpha_k^6 + 2\alpha_k^5 + 3\alpha_k^4 + 4\alpha_k^3 + 5\alpha_k^2 + 6\alpha_k + 7},$$

$$Q_m^* = \sum_{k=1}^7 \alpha_k^m \quad \text{and}$$

$$E_m^* = \sum_{k=1}^7 \frac{(\alpha_k - 1)\alpha_k^{m+5}}{2\alpha_k^6 + 2\alpha_k^5 + 3\alpha_k^4 + 4\alpha_k^3 + 5\alpha_k^2 + 6\alpha_k + 7}$$

respectively.

**3.6 Theorem 2.**

For  $r, n \in \mathbb{N}$ , Catalan's identity holds as;

$$E_{r+n}^* E_{r-n}^* - E_r^{*2} = (P_{r+n}^* - P_{r+n-1}^*)(P_{r-n}^* - P_{r-n-1}^*) - (P_r^* - P_{r-1}^*)^2$$

For  $n = 1$  we have Cassini identity as;

$$E_{r+1}^* E_{r-1}^* - E_r^{*2} = (P_{r+1}^* - P_r^*)(P_{r-1}^* - P_{r-2}^*) - (P_r^* - P_{r-1}^*)^2$$

**3.7 Theorem 3.**

The following identities are true for  $r, n \in \mathbb{Z}$ :

(i): (Melham's identity)

$$E_{r+1}^* E_{r+2}^* E_{r+6}^* - E_{r+3}^{*3} = (P_{r+1}^* - P_r^*)(P_{r+2}^* - P_{r+1}^*)(P_{r+6}^* - P_{r+5}^*)(P_{r+1}^* - P_r^*) - (P_{r+3}^* - P_{r+2}^*)^3$$

(ii): (Gelin-Cesaro's identity)

$$E_{r+2}^* E_{r+1}^* E_{r-1}^* E_{r-2}^* - E_r^{*6} = (P_{r+2}^* - P_{r+1}^*)(P_{r+1}^* - P_r^*)(P_{r-1}^* - P_{r-2}^*)(P_{r-2}^* - P_{r-3}^*) - (P_r^* - P_{r-1}^*)^4$$

(iii): (d'Ocagnes identity)

$$E_{n+1}^* E_r^* - E_n^* E_{r+1}^* = (P_{n+1}^* - P_n^*)(P_r^* - P_{r-1}^*) - (P_n^* - P_{n-1}^*)(P_{r+1}^* - P_r^*)$$

**4. Linear sum of pell, pell-like and modified pell sequences**

**4.1 Theorem**

For  $m \geq 0$  the sum of the generalized seventh order Pell numbers we have:

$$\sum_{k=0}^m H_k = \frac{1}{7}(H_{m+7} - H_{m+6} - 2H_{m+5} - 3H_{m+4} - 4H_{m+3} - 5H_{m+2} - 6H_{m+1} - H_6 + H_5 - 2H_4 - 3H_3 + 4H_2 + 5H_1 + 6H_0)$$

Proof. : Using the recurrence relation, then, solving the above equality we obtain

$$H_m = 2H_{m-1} + H_{m-2} + H_{m-3} + H_{m-4} + H_{m-5} + H_{m-6} + H_{m-7}$$

i.e.,

$$H_{m-7} = H_m - 2H_{m-1} - H_{m-2} - H_{m-3} - H_{m-4} - H_{m-5} - H_{m-6}$$

For  $m = 7, 8, 9, 10, \dots$  we have the pattern as;

$$H_0 = H_7 - 2H_6 - H_5 - H_4 - H_3 - H_2 - H_1,$$

$$H_1 = H_8 - 2H_7 - H_6 - H_5 - H_4 - H_3 - H_2,$$

$$H_2 = H_9 - 2H_8 - H_7 - H_6 - H_5 - H_4 - H_3,$$

$$\dots = \dots$$

$$H_{m-7} = H_m - 2H_{m-1} - H_{m-2} - H_{m-3} - H_{m-4} - H_{m-5} - H_{m-6}$$

$$H_{m-6} = H_{m+1} - 2H_m - H_{m-1} - H_{m-2} - H_{m-3} - H_{m-4} - H_{m-5}$$

$$H_{m-5} = H_{m+2} - 2H_{m+1} - H_m - H_{m-1} - H_{m-2} - H_{m-3} - H_{m-4}$$

$$H_{m-4} = H_{m+3} - 2H_{m+2} - H_{m+1} - H_m - H_{m-1} - H_{m-2} - H_{m-3}$$

$$H_{m-3} = H_{m+4} - 2H_{m+3} - H_{m+2} - H_{m+1} - H_m - H_{m-1} - H_{m-2}$$

$$H_{m-2} = H_{m+5} - 2H_{m+4} - H_{m+3} - H_{m+2} - H_{m+1} - H_m - H_{m-1}$$

$$H_{m-1} = H_{m+6} - 2H_{m+5} - H_{m+4} - H_{m+3} - H_{m+2} - H_{m+1} - H_m$$

$$H_m = H_{m+7} - 2H_{m+6} - H_{m+5} - H_{m+4} - H_{m+3} - H_{m+2} - H_{m+1}$$

Now, by rearrangement we have

$$\sum_{k=0}^m H_k = (H_{m+7} + H_{m+6} + H_{m+5} + H_{m+4} + H_{m+3} + H_{m+2} + H_{m+1} - H_6 - H_5 - H_4 - H_3 - H_2 - H_1 - H_0 + \sum_{k=0}^m H_k) - 2(H_{m+6} + H_{m+5} + H_{m+4} + H_{m+3} + H_{m+2} + H_{m+1} - H_5 - H_4 - H_3 - H_2 - H_1 - H_0 + \sum_{k=0}^m H_k) - (H_{m+5} + H_{m+4} + H_{m+3} + H_{m+2} + H_{m+1} - H_4 - H_3 - H_2 - H_1 - H_0 + \sum_{k=0}^m H_k) - (H_{m+4} + H_{m+3} + H_{m+2} + H_{m+1} - H_3 - H_2 - H_1 - H_0 + \sum_{k=0}^m H_k) - (H_{m+3} + H_{m+2} + H_{m+1} - H_2 - H_1 - H_0 + \sum_{k=0}^m H_k) - (H_{m+2} + H_{m+1} - H_1 - H_0 + \sum_{k=0}^m H_k) - (H_{m+1} - H_0 + \sum_{k=0}^m H_k)$$

**4.2 Corollary 7.**

For  $m \geq 0$  the sum of the seventh order  $\{P_m^*\}_{m \geq 0}$ ,

$\{Q_m^*\}_{m \geq 0}$  and  $\{E_m^*\}_{m \geq 0}$  sequences are

(a) 
$$\sum_{k=0}^m P_k^* = \frac{1}{7}(P_{k+7}^* - P_{k+6}^* - 2P_{k+5}^* - 3P_{k+4}^* - 4P_{k+3}^* - 5P_{k+2}^* - 6P_{k+1}^* - 1)$$

(b) 
$$\sum_{k=0}^m Q_k^* = \frac{1}{7}(Q_{k+7}^* - Q_{k+6}^* - 2Q_{k+5}^* - 3Q_{k+4}^* - 4Q_{k+3}^* - 5Q_{k+2}^* - 6Q_{k+1}^* - 1)$$

(c) 
$$\sum_{k=0}^m E_k^* = \frac{1}{7}(E_{k+7}^* - E_{k+6}^* - 2E_{k+5}^* - 3E_{k+4}^* - 4E_{k+3}^* - 5E_{k+2}^* - 6E_{k+1}^* - 1)$$

**5. Conclusions**

After derived an explicit formula for the generalized pell sequences and pell-like sequences of order seven. These derivations may extend higher order of generalized pell

sequences and pell-like sequences for further expansion to innovative ways.

### Acknowledgment

The author is indebted to the worthy supervisor for her valuable comments, suggestions, additional references and further improving the presentation of this paper. We are also equally thankful to the CT University Punjab for providing necessary resources to conduct this research.

### References

- [1]. NA Bicknell. Primer on the Pell sequence and related sequence, *Fibonacci Quarterly*, 13(4), 1975, 345-349.
- [2]. A Dasdemir. On the Pell, Pell-Lucas and Modified Pell Numbers by Matrix Method, *Applied Mathematical Sciences*, 5(64), 2011, 3173-3181.
- [3]. J Ercolano. Matrix generator of Pell sequence, *Fibonacci Quarterly*, 17(1), 1979, 71-77.
- [4]. H Gökbas, H. Köse. Some sum formulas for products of Pell and Pell-Lucas numbers, *Int. J. Adv. Appl. Math. and Mech.* 4(4), 2017, 1-4.
- [5]. CA Hanusa. A Generalized Binet's Formula for kth Order Linear Recurrences: A Markov Chain Approach, Harvey Mudd College, Undergraduate Thesis (Math Senior Thesis), 2001.
- [6]. AF Horadam. Pell identities, *Fibonacci Quarterly*, 9(3), 1971, 245-263.
- [7]. D Kalman. Generalized Fibonacci Numbers by Matrix Methods, *Fibonacci Quarterly*, 20(1), 1982, 73-76, 1982.
- [8]. E Kiliç, D Ta,Şçi. The Linear Algebra of The Pell Matrix, *Boletín de la Sociedad Matemática Mexicana*, 3(11), 2005.
- [9]. E Kiliç, D Ta,Şçi. The Generalized Binet Formula, Representation and Sums of the Generalized Order-k Pell Numbers, *Taiwanese Journal of Mathematics*, 10(6), 2006, 1661-1670.
- [10]. E Kiliç, P Stanica. A Matrix Approach for General Higher Order Linear Recurrences, *Bulletin of the Malaysian Mathematical Sciences Society*, (2) 34(1), 2011, 51-67.
- [11]. T Koshy. *Pell and Pell-Lucas Numbers with Applications*, Springer, New York, 2014
- [12]. R Melham. Sums Involving Fibonacci and Pell Numbers, *Portugaliae Mathematica*, 56(3), 1999, 09-317.
- [13]. NJA Sloane, The on-line encyclopedia of integer sequences, <http://oeis.org/>
- [14]. N Ahmad, L Prasher. *Vidyabharati International Interdisciplinary Research Journal Generalized Formulae for Lucas-like Sequence of 6th to 9th Order.*
- [15]. N Ahmad, L Prasher. Findings for Generalized Sixth, Seventh, Eighth and Ninth order of Fibonacci-Like Sequences.
- [16]. *Vidyabharati International Interdisciplinary Research Journal*,
- [17]. Y Soykan. On Generalized Third-Order Pell Numbers, *Asian Journal of Advanced Research and Reports*, 6(1), 2019, 1-18.
- [18]. Y Soykan. A Study of Generalized Fourth-Order Pell Sequences, *Journal of Scientific Research and Reports*, 25(1-2), 2019, 1-18.
- [19]. Y Soykan. Properties of Generalized Fifth-Order Pell Numbers, *Asian Research Journal of Mathematics*, 15(3), 2019, 1-18.
- [20]. T Yagmur. New Approach to Pell and Pell-Lucas Sequences, *Kyungpook Math. J.* 59, 219, 23-34, 2019.
- [21]. S Yüksel. On Generalized Sixth-Order Pell Sequences. *Journal of Scientific Perspectives* 4.1, 2020, 49-70.
- [22]. S Vajda, *Fibonacci and Lucas numbers and the golden section. Theory and Applications*, Ellis Horwood Limited, 1989.
- [23]. J Bravo, L Jose. Herrera, F Luca. On a generalization of the Pell sequence. *Mathematica Bohemica* 146 (2), 2021, 199-213.
- [24]. C Paula, P Vasco. Modified k-Pell sequence: Some identities and ordinary generating function. *Appl. Math. Sci* 121 (7), 2013, 6031-6037.